



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 433 900 A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90124133.1

(51) Int. Cl. 5: **C12N 15/12**, C07K 13/00,
C12N 15/79

(22) Date of filing: 13.12.90

(30) Priority: 13.12.89 IL 92697
12.07.90 IL 95064

(43) Date of publication of application:
26.06.91 Bulletin 91/26

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: **YEDA RESEARCH AND
DEVELOPMENT COMPANY LIMITED**
P.O. Box 95
Rehovot 76100(IL)

(72) Inventor: Wallach, David
24, Borochoy Street
Rehovot 76100(IL)
Inventor: Nophar, Yaron
56, Bialic Street
Ramat Gan, 52441(IL)
Inventor: Kemper, Oliver
Anemonenweg 10
W-6719 Bockenheim(DE)
Inventor: Engelmann, Hartmut
Joseph-Lutz-Weg 35
W-8000 München 70(DE)
Inventor: Brakebusch, Cord
Salzbahlumer Weg 9
W-3300 Braunschweig(DE)
Inventor: Aderka, Dan
4, Avivim Street
Holon(IL)

(74) Representative: Vossius & Partner
Siebertstrasse 4 P.O. Box 86 07 67
W-8000 München 86(DE)

(94) Expression of the recombinant tumor necrosis factor binding protein I (TBP-I).

(57) Tumor Necrosis Factor Binding Protein I (TBP-I), precursors and analogs thereof, are produced by transfecting eukaryotic cells with an expression vector comprising a DNA molecule encoding the whole human type I TNF receptor or a soluble domain thereof, and culturing the transfected cells, whereby the soluble proteins are secreted into the medium.

EP 0 433 900 A1

EXPRESSION OF THE RECOMBINANT TUMOR NECROSIS FACTOR BINDING PROTEIN I (TBP-I)

The present invention relates to human Tumor Necrosis Factor (TNF) Binding Protein I, herein designated TBP-I, and more particularly, to the cloning of the gene coding for said protein and its expression in host cells.

TNF- α and TNF- β (lymphotoxin) are structurally related polypeptide cytokines, produced primarily by mononuclear leukocytes, whose effects on cell function constitute a major factor in the elicitation of the inflammatory response. The TNFs affect cells in different ways, some of which resemble the functional modes of other inflammatory mediators, like interleukin-1 (IL-1) and interleukin-6 (IL-6). What appears most distinctive regarding the activity of the TNFs is that many of their effects can result in cell and tissue destruction. Increasing evidence that over-induction of these destructive activities contributes to the pathogenesis of a number of diseases, makes it of particular interest to elucidate their mechanisms and the ways they are regulated (Old, L.J. (1988) *Sci.Am.* 258, pp. 41-49).

High affinity receptors, to which both TNF- α and TNF- β bind (Beutler, B.A., et al. (1985) *J.Exp.Med.* 161, pp. 984-995) play a key role in the initiation and in the control of the cellular response to these cytokines. These receptors are expressed on the surfaces of a variety of different cells. Studies showing that antibodies reacting with their extracellular portions affect cells in a manner very similar to the TNFs, demonstrate that the receptors and cellular components associated with them are sufficient to provide the intracellular signalling for the effects of the TNFs (Espevik, T., et al., (1990) *J.Exp.Med.* 171, pp. 415-426).

Other studies have shown that molecules related to the TNF receptors (TNF-Rs) exist also in soluble forms. Two immunologically distinct species of such soluble TNF-Rs, designated TNF Binding Proteins I and II, or TBP-I and TBP-II, respectively, were recently isolated from human urine (Engelmann, H., et al., (1989) *J.Biol.Chem.* 264, pp. 11974-11980; Engelmann, H., et al., (1990) *J.Biol.Chem.* 265, pp. 1531-1536; Olsson, I., et al., (1989) *Eur.J.Haematol.* 42, pp. 270-275; Seckinger, P., et al., (1989a) *J.Biol.Chem.* 264, pp. 11966-11973). Immunological evidence indicated that the two proteins are structurally related to two molecular species of the cell surface TNF-R (the type I and type II receptors, respectively). Antibodies to each of the two soluble proteins were shown to block specifically the binding of TNF to one of the two receptors and could be used to immunoprecipitate the receptors. Antibodies against one of the two soluble proteins (TBP-I) were also found to induce effects characteristic of TNF in cells which express the immunologically cross-reactive cell receptors (Engelmann, H., et al., (1990) *ibid.*). Like the cell surface receptors for TNF, the soluble forms of these receptors specifically bind TNF and can thus interfere with its binding to cells. It was suggested that they function as physiological inhibitors of TNF activity (Engelmann et al., 1989 (*ibid.*); Olsson et al., 1989 (*ibid.*); Seckinger et al., 1989a (*ibid.*)).

Soluble forms of cell surface receptors may be derived from the cell surface form of the receptor by proteolytic cleavage, or by a different mechanism proposed in two recent studies describing the cloning of the cDNAs for the receptors to IL-4 and IL-7. Besides cDNA clones encoding the full length receptors, clones which encode truncated, soluble forms of these receptors were also isolated in these studies. It was suggested that these latter clones are derived from transcripts specifically encoding soluble forms of the receptors, transcribed from the same genes which encode the cell surface forms, but differently spliced (Mosley, B., et al., (1989) *Cell* 59, pp. 335-348; Goodwin, R.G., et al., (1990) *Cell* 60, pp. 941-951).

Two recent studies have described the molecular cloning and expression of human type I TNF cell surface receptor (Loetscher, H., et al. (1990) *Cell* 61, pp. 351-359; Schall, T.J., et al., (1990) *Cell* 61, pp. 361-370).

The present invention relates to the production of human TBP-I, precursors and analogs thereof, by a method comprising transfection of eukaryotic, preferably CHO cells, with an expression vector comprising a DNA molecule encoding the whole type I human TNF receptor or a soluble domain thereof. When the whole DNA molecule is used, soluble proteins are produced by the transfected cells, along with the cell surface receptor, and are secreted into the medium.

The invention further relates to soluble proteins selected from precursors and analogs of TBP-I, which are secreted into the medium by eukaryotic cells transfected with a DNA molecule encoding the whole human type I TNF receptor or a soluble domain thereof.

Figure 1 shows the nucleotide sequence of the type I TNF receptor cDNA and the predicted amino acid sequence of the encoded protein.

(A) shows the probes used for screening for the cDNA, wherein:

(a) shows the NH₂-terminal amino acid sequence of TBP-I;

(b) shows synthetic oligonucleotide probes, designed on the basis of the NH₂-terminal amino acid sequence, used for initial screening; and

(c) and (d) are probes overlapping with (b), used to confirm the validity of clones isolated in the initial screening.

(B) is the schematic presentation of the cDNA clones isolated from a human colon (C2) and from CEM-lymphocytes (E13) libraries and a diagram of the complete cDNA structure. Untranslated sequences are represented by a line. Coding regions are boxed. The shaded portions represent the sequences which encode the signal peptide and the transmembrane domains.

(C) shows the hydropathy profile of the predicted amino acid sequence of the TNF receptor. Hydrophobicity (above the line) and hydrophilicity (below the line) values were determined using the sequence analysis software package of the University of Wisconsin genetic computer group (UWCG) according to Kyte and Doolittle (1982). The curve is the average of the hydrophobicity index for each residue over a window of nine residues.

(D) depicts the nucleotide and predicted amino acid sequences of the type I TNF receptor. The presumptive start and stop signals are denoted by asterisks; the three sequences derived from TBP-I by broken overlining; the transmembrane and leader domains by round-ended boxes; and the four repetitive sequences in the extracellular domain by thick underlining. Cysteine residues are boxed. Glycosylation sites are overlined and the presumptive polyadenylation signal is underlined.

Figure 2 shows the detection of type I TNF-R using monoclonal antibodies to TBP-I in CHO cells transfected with E13 cDNA. CHO cells, clones R-18 (transfected with an expression vector in which the E13 cDNA was placed under the control of an SV40 promoter) and C-6 (control; a clone of cells transfected with an expression vector in which E13 was placed in the inverse orientation), and HeLa cells, were stained with the anti-TBP-I monoclonal antibodies 17, 18, 20 and 30 followed by incubation with FITC conjugated anti-mouse F(ab). Fluorescence intensity was compared with that observed when a mouse monoclonal antibody against TNF was used in the first step of the staining as a control.

Figure 3 shows reversed phase HPLC of the CHO-produced, soluble form of the type I TNF-R.

A concentrate of the conditioned medium of the CHO R-18 clones (see Fig. 2) and a concentrate of the CHO C-6 clone to which 3 ng pure TBP-I was added, were applied to an Aquapore RP300 column. Elution was performed with a gradient of acetonitrile in 0.3% aqueous trifluoroacetic acid (---). Fractions were examined for content of protein (—) and of the soluble form of the type I receptor by an ELISA

(---),

(as described in Example 1: Procedures). None of the eluted fractions of a concentrate of the CHO C-6 clone, without addition of TBP-I, was found to contain any detectable amounts of the soluble form of the receptor (not shown).

Figure 4 shows the time course of the release of COOH-terminal amino acids from TBP-I by carboxypeptidase Y.

Figure 5 shows the construction of plasmid pSV-TBP, which contains the DNA sequence encoding TBP-I fused to the strong SV40 early gene promoter.

Figure 6 shows the construction of the plasmid pCMV-TBP, which contains the DNA sequence encoding TBP-I fused to the cytomegalovirus (CMV) promoter.

Purified TBP-I isolated from human urine was described in European Patent Application EP 0 308 378 of the present applicants and shown to contain at the N-terminus the amino acid sequence shown in Fig. 1Aa.

The COOH-terminal of TBP-I was determined now and shown to contain a major fraction containing the sequence Ile-Glu-Asn denoted by broken overlining at positions 178-180 in Fig. 1D, and at least one minor fraction including at least two further amino acids Val-Lys at positions 181-182.

The invention relates to a method for the production of a soluble recombinant protein selected from human Tumor Necrosis Factor Binding Protein I (TBP-I), biologically active precursors and analogs thereof, which comprises:

- i) transfecting eukaryotic cells with an expression vector comprising a DNA molecule encoding the whole type I human TNF receptor or a soluble domain thereof, and
- ii) culturing the transfected cells, whereby the desired protein is produced and secreted into the medium.

The DNA sequence encoding the whole type I TNF receptor is depicted in Figure 1D. The soluble domain thereof includes the sequence down to position 180 (Asn) or 182 (Lys) or even some additional amino acids after position 182.

The soluble proteins produced by the transfected cells according to the method of the invention and

secreted into the medium may have at the N-terminus the sequence Asp-Ser-Val denoted by broken overlining at positions 20-23 in Fig. 1D (TBP-I), or the sequence Leu-Val-Pro at positions 9-11 or Ile-Tyr-Pro at positions 1-3 or any other sequence between Ile(+1) and Asp(20). The proteins may have at the COOH terminal any of the sequences described above. All these soluble proteins, if biologically active with TBP-I-
 5 like activity, are encompassed by the invention as precursors and analogs of TBP-I.

According to the invention, oligonucleotide probes designed on the basis of the NH₂-terminal amino acid sequence of TBP-I, were synthesized by known methods and used for screening for the cDNA coding for TBP-I in cDNA libraries. In a human colon cDNA library, a C2 cDNA insert was found which hybridized to said probes and it was used for further screening in a human CEM-lymphocytes lambda ZAP cDNA
 10 library, thus leading to the cDNA shown in Fig. 1D.

The DNAs of positive clones were then inserted into appropriately constructed expression vectors by techniques well known in the art. In order to be capable of expressing a desired protein, an expression vector should comprise also specific nucleotide sequences containing transcriptional and translational regulatory information linked to the DNA coding for the desired protein in such a way as to permit gene
 15 expression and production of the protein. The gene must be preceded by a promoter in order to be transcribed. There are a variety of such promoters in use, which work with different efficiencies (strong and weak promoters).

The DNA molecule comprising the nucleotide sequence coding for a protein comprising the amino acid sequence of TBP-I, i.e. TBP-I, a precursor or an analog thereof, preceded by a nucleotide sequence of a signal peptide and the operably linked transcriptional and translational regulatory signals is inserted into a
 20 vector which is capable of integrating the desired gene sequences into the host cell chromosome. The cells which have stably integrated the introduced DNA into their chromosomes can be selected by also introducing one or more markers which allow for selection of host cells which contain the expression vector.

In a preferred embodiment, the introduced DNA molecule will be incorporated into a plasmid or viral
 25 vector capable of autonomous replication in the recipient host. Factors of importance in selecting a particular plasmid or viral vector include the ease with which recipient cells that contain the vector may be recognized and selected from those recipient cells which do not contain the vector; the number of copies of the vector which are desired in a particular host and whether it is desirable to be able to "shuttle" the vector between host cells of different species. Once the vector or DNA sequence containing the construct(s) has
 30 been prepared for expression, the DNA construct(s) may be introduced into an appropriate host cell by any of a variety of suitable means: transformation, transfection, conjugation, protoplast fusion, electroporation, calcium phosphate precipitation, direct microinjection, etc.

Host cells to be used in this invention may be either prokaryotic or eukaryotic. Prokaryotic hosts, such as bacteria, e.g. E.coli, are used only when the cDNA encoding the soluble domain of the type I TNF
 35 receptor is used to transfect the cells. Under such conditions, the protein will not be glycosylated. The prokaryotic host must be compatible with the replicon and control sequences in the expression plasmid.

Eukaryotic cells are transfected according to the invention with plasmids comprising the cDNA encoding the whole type I TNF receptor. Preferred eukaryotic hosts are mammalian cells, e.g., human, monkey, mouse and chinese hamster ovary (CHO) cells. They provide the soluble form of the protein, besides the
 40 cell surface receptor, and provide post-translational modifications to protein molecules including correct folding or glycosylation at correct sites. The eukaryotic cells may also be transfected with a plasmid comprising a cDNA encoding a soluble domain of the human type I TNF receptor molecule. Preferred mammalian cells according to the invention are CHO cells.

After the introduction of the vector, the host cells are grown in a selective medium, which selects for the
 45 growth of vector-containing cells. Expression of the cloned gene sequence(s) results in the production of the desired soluble protein, that is secreted into the medium, and may then be isolated and purified by any conventional procedure involving extraction, precipitation, chromatography, electrophoresis, or the like.

In a preferred embodiment, CHO cells are transfected with the type I TNF-R cDNA shown in Fig. 1D and they produce both the cell surface receptor and TBP-I, its soluble form, and/or precursors and analogs
 50 thereof.

The data presented in the present application are consistent with the notion that TBP-I - the soluble form for the type I TNF-R - constitutes a fragment of the cell surface form of this receptor, corresponding to its extracellular domain. The receptor is recognized by several monoclonal antibodies to TBP-I which
 55 interact with several spatially distinct epitopes in this protein. The amino acid sequence in the extracellular domain matches the sequence of TBP-I.

Particularly informative with regard to the mechanism of formation of TBP-I is the finding described in the present application, that a soluble form of the type I TNF-R is produced by CHO cells which were transfected with the TNF-R cDNA. This implies that cells possess some mechanism(s) which allow(s) the

formation of the soluble form of the TNF-R from that same transcript that encodes the cell surface form.

The low rate of production of the soluble form of the type I TNF-R by the E13-transfected CHO cells does not necessarily reflect maximal activity. In HT29 cells, the spontaneous release of a soluble form of type I TNF-R occurs at about a 10-fold higher rate than that observed with the CHO-R-18 clone.

5 A likely mechanism whereby soluble forms of TNF receptors can be derived from the same transcripts which encode the cell surface forms is proteolytic cleavage. Indeed, flanking the amino acid residue which corresponds to the NH₂-terminus of TBP-I there are, within the sequence of amino acids of the receptor, two basic amino acid residues (Lys-Arg) which can serve as a site of cleavage by trypsin-like proteases. The identity of the proteases which might cause cleavage to take place at the COOH terminus of TBP-I is
10 not known.

The invention will be illustrated by the following examples:

EXAMPLE 1: PROCEDURES

15 A) Determination of amino acid sequences within the TNF-binding proteins TBP-I and TBP-II

The TNF Binding Proteins TBP-I and TBP-II were isolated from concentrated preparations of urinary proteins, as described previously (Engelmann, H., et al., (1990) J.Biol.Chem. 265, pp. 1531-1536) by ligand (TNF) affinity chromatography followed by reversed phase HPLC. TBP-I was cleaved with cyanogen
20 bromide, yielding two peptides which, following reduction and alkylation, were isolated by reversed phase HPLC. The two peptides (CNBr-1 and CNBr-2 in Table I) were subjected to NH₂-terminal sequence analysis or a pulsed liquid gas phase protein microsequencer (Model 475A, Applied Biosystems Inc., Foster City CA). The sequence found for one of the peptides (CNBr-1) was identical to the NH₂ sequence of the intact TBP-I protein.

25 The COOH terminal sequence of amino acids in TBP-I was determined by digestion of the protein with carboxypeptidase Y followed by sequential analysis of the released amino acids. A sample of pure TBP-I (32 µg) was mixed with 1 nmole of norleucine, as an internal standard, dried thoroughly and resuspended in 8 µl 0.1 M sodium acetate buffer, pH-5.5, containing 0.8 µg carboxypeptidase Y (Sigma, St. Louis, MO). Digestion was performed at room temperature, 2 µl Aliquots withdrawn at various time points were acidified
30 by adding 3 µl of 10% acetic acid to each, followed by addition of 15 µl 0.5% EDTA. They were then subjected to automated amino acid analysis (Applied Biosystems, U.K. model 420A). The results (shown in Fig. 4) indicate the sequence -Ile-Glu-Asn-COOH. Minor fractions were detected containing two or more additional amino acids.

Sequences within TBP-II were determined by generation of tryptic peptides of the protein. A sample of
35 pure TBP-II (200 µg) was reduced, alkylated and repurified on an Aquapore RP-300 reversed-phase HPLC column. Fractions containing the modified protein were pooled and the pH was adjusted to 8.0 with NaHCO₃. Digestion with TPCK-trypsin (238 U/mg, Millipore Corp., Freehold, NJ) was performed for 16 h. at room temperature at an enzyme to substrate ratio of 1:20 (w/w). The digest was loaded on a C₁₈ RP-P reversed phase HPLC column (Synchrom, Linden, IN) and the peptides separated by a linear 0 to 40%
40 acetonitrile gradient in 0.3% aqueous trifluoroacetic acid. The NH₂ terminal amino acid sequences of the peptides and of the intact protein (N-terminus) are presented in Table I. The peptides were numbered according to their sequences of elution from the RP-P column. In the fractions denoted as 39,44,46,53 and 54, where heterogeneity of sequences was observed, both the major and the secondary sequences are presented.

45 b) Isolation of cDNA clones

Three mixtures of synthetic oligonucleotide probes (Figs. 1Ab, 1Ac) generated from the nucleotide sequence deduced from the NH₂-terminal amino acid sequence of TBP-I (Fig. 1Aa) were used for the
50 screening of cDNA libraries. Initial screenings were carried out with 48-fold degenerated, 26-mers into which deoxyinosine was introduced, wherever the codon ambiguity allowed for all four nucleotides (Fig. 1Ab). The validity of positive clones was examined by testing their hybridization to two mixed 17-mer nucleotide sequences containing 96 and 128 degeneracies, corresponding to two overlapping amino acid sequences which constitute part of the sequences to which the 26-base probes correspond (Fig. 1Ac and d). An
55 oligonucleotide probe corresponding to a sequence located close to the 5' terminus of the longest of the partial cDNA clones isolated with the degenerated probes (nucleotides 478-458 in Fig. 1D) was applied for further screening cDNA libraries for a full length cDNA clone. ³²P-labeling of the probes, using T4 polynucleotide kinase, plating of the phages in lawns of bacteria, their screening with the radio-labelled

probes, isolation of the positive clones and subcloning of their cDNA inserts were carried out using standard procedures (Sambrook, J., et al., (1989) Molecular Cloning, A Laboratory Manual. Cold Spring Harbor Laboratory Press).

5 c) Nucleotide sequencing of the cDNA clones

cDNA inserts isolated from positive lambda GT11 recombinant phages were subcloned into the pBluescript KS(-) vector. Inserts found in lambda ZAP phages were rescued by excising the plasmid pBluescript SK(-) in them, using the R408 helper phage (Short, J.M., et al., (1988) Nucl.Acids Res. 16, pp. 7583-7600). DNA sequencing in both directions was done by the dideoxy chain termination method. Overlapping deletion clones of the cDNAs were generated, in both orientations, by digestion of the cDNA with exonuclease III ("Erase a base" kit, Promega Biotec, Madison, WI). Single stranded templates derived from these clones using the R408 phage were sequenced with a T7 DNA polymerase sequencing system (Promega).

15 d) Constitutive expression of the type I human TNF-R in CHO cells

The E13 insert was introduced into a modified version of the pSVL expression vector. This construct was transfected, together with the pSV2-DHFR plasmid which contains the DHFR cDNA, into DHFR deficient CHO cells, using the calcium phosphate precipitation method. Transfection with a recombinant pSVL vector which contained the E13 insert in the inverse orientation served as a control. Cells expressing the DHFR gene were selected by growth in nucleotide-free MEM alpha medium containing fetal calf serum which had been dialyzed against phosphate buffered saline. Individual clones were picked out and then further selected for amplification of the transfected cDNAs by growth in the presence of 500 nM sodium methotrexate.

e) Detection of surface-expressed type I TNF-R in the CHO cells

Binding of radiolabelled human rTNF to cells (seeded in 15 mm tissue culture plates at a density of 2.5×10^5 cells/plate) was quantitated as described before (Holtmann, H. and Wallach, D. (1987) J.Immunol. 139, pp. 1161-1167).

To examine the binding of monoclonal antibodies against TBP-I to CHO cells, the cells were detached by incubation in phosphate buffered saline (PBS: 140 mM NaCl, 1.5 mM KH_2PO_4 , 8 mM Na_2HPO_4 , 2.7 mM KCl, 0.5 mM MgCl_2 , 0.9 mM CaCl_2), containing 5 mM EDTA and then incubated for 45 min. at 0°C with 50 $\mu\text{g/ml}$ of the test monoclonal antibody in PBS containing 0.5% bovine serum albumin, and 15 mM sodium azide (PBS/BSA). After washing the cells with PBS/BSA they were incubated further for 30 min. at 0°C with FITC labelled, affinity purified goat antibody to the F(ab) fragment of mouse IgG (1:20 in PBS/BSA) (Bio-Makor, Israel) and then analyzed by determining the intensity of fluorescence in samples of 10^4 cells using the Becton Dickinson fluorescence activated cell sorter 440. Three monoclonal antibodies to TBP-I, clones 17,18 and 20, shown by cross competition analysis to recognize four spatially distinct epitopes in the TBP-I molecule (European Patent Application No. 90115105.0) and, as a control, a monoclonal antibody against TNF- α (all purified from ascitic fluids by ammonium sulphate precipitation and of the IgG2 isotype), were used.

45 f) Quantitation of the soluble form of the type I TNF-R by ELISA

A sensitive enzyme linked immunosorbent assay was set up using TBP-I-specific monoclonal and polyclonal antibodies in a sandwich technique. Immunoglobulins of the anti-TBP-I mAb clone 20 (European Patent Application No. 90115105.0) were adsorbed to 96-well ELISA plates (maxisorp, Nunc, Denmark) by incubation of the plates for 2 h. at 37°C with a solution of 25 $\mu\text{g/ml}$ of the antibody in PBS. After incubating the wells further for 2 h. at 37°C with a solution containing phosphate buffered saline, 1% BSA, 0.02% NaN_3 and 0.05% Tween 20 (blocking solution) to block nonspecific further binding of protein, tested samples were applied in aliquots of 50 $\mu\text{l/well}$. The plates were then incubated for 2 h. at 37°C , rinsed 3 times with PBS supplemented with 0.05% Tween 20 (washing solution) and rabbit polyclonal antiserum against TBP-I, diluted 1:500 in blocking solution, was added to the wells. After further incubation for 12 h. at 4°C the plates were rinsed again and incubated for 2 h. with horse raddish peroxidase-conjugated purified goat anti-rabbit IgG. The assay was developed using 2,2'-azino-bis (3-ethylbenzthiazoline-6 sulfonic acid) as a substrate (Sigma). The enzymatic product was determined colorimetrically at 600 nm. Pure TBP-I served

as a standard.

g) Detection of a soluble form of the type I TNF-R in the growth medium of the transfected CHO cells and its analysis by reversed phase HPLC

5 The amounts of the soluble form of the type I TNF-R in samples of the medium of the tested CHO cells, collected 48 h after medium replacement, were determined by the immunoassay described above. For analysis of the soluble receptor by reversed phase HPLC the CHO cells were cultured for 48 h. in serum-free medium (nucleotide-free MEM α). The medium samples were concentrated 100-fold by ultrafil-
10 tration on an Amicon PM5 membrane and 100 μ l aliquots were then applied to an Aquapore RP300 column (4.5 X 30 mm, Brownlee Labs) preequilibrated with 0.3% aqueous trifluoroacetic acid. The column was washed with this solution at a flow rate of 0.5 ml/min until all unbound proteins were removed, and then eluted with a gradient of acetonitrile concentration in 0.3% aqueous trifluoroacetic acid, as described before (Engelmann, H., et al., (1989) J.Biol.Chem. 264, pp. 11974-11980). Fractions of 0.5 ml were collected and,
15 after concentration in vacuo, were neutralized with 1 M HEPES buffer pH 9.0. Amounts of soluble type I TNF-R in the fractions were determined by ELISA and the concentration of protein by the fluorescamine method.

EXAMPLE 2

a) Cloning of the cDNA for the Type I TNF-R

To clone the cDNAs which code for the TNF-binding protein, TBP-I, and its related TNF receptor, several cDNA libraries were screened, using 3 overlapping oligonucleotide probes designed on the basis of
25 the NH₂-terminal amino acid sequence of TBP-I (Fig. 1A). In a lambda GT11 library derived from the mRNA of human colon (randomly primed, Clontech, Palo Alto, CA), four recombinant phages which hybridized with the three probes were detected. The inserts in these four phages were similar in size, and were found to overlap by restriction mapping and sequence analysis.

Complete analysis of the sequence of the longest of the four (C2 in Fig. 1B, deposited on 6.12.1989
30 with the Collection Nationale de Cultures de Microorganismes (C.N.C.M.), Paris, France, Accession No. I-917) revealed an open reading frame, extended over its entire length. A polypeptide chain encoded in this reading frame fully matches the NH₂-terminal amino acid sequence of TBP-I. Neither an initiation nor a stop codon was found in the C2 insert. Rescreening the colon cDNA library, using another probe corresponding to a sequence found in C2 (see Example 1: Procedures), yielded several other recombinant phages
35 containing inserts that overlap with the C2 insert. However, none of them provided further sequence information on the cDNA in the 5' or the 3' direction. In a lambda ZAP cDNA library derived from the mRNA of CEM lymphocytes (Oligo dT and randomly primed, Clontech) five phages hybridizing with this probe were detected, which contained significantly longer inserts than C2.

The longest insert (E13, Fig. 1B) was sequenced in its entirety (Fig. 1D) and was found to contain the
40 C2 sequence (nucleotides 346-1277 in Fig. 1D) within one long open reading frame of 1365 bp, flanked by untranslated regions of 255 and 555 nucleotides at its 5' and 3' ends, respectively. The potential ATG initiation site, occurring at positions 256-258 in the nucleotide sequence (denoted by an asterisk in Fig. 1D) is preceded by an upstream in-frame termination codon at bases 244-246. The start location is in conformity with one of the possible alternatives for the translation initiation consensus sequence
45 (GGCATGG, nucleotides 253-259).

There is no characteristic poly(A) addition signal near the 3' end of the cDNA. The sequence ACTAAA, at nucleotides 2045-2050, may serve as an alternative to this signal, but with low efficiency. At nucleotides 1965-2000, there are six consecutive repeats of the sequence G(T)_n (n varying between 4 and 8).

The size of the protein encoded by the cDNA (about 50 kD) is significantly larger than that of TBP-I. A
50 hydropathy index computation of the deduced amino acid sequence of the protein (Fig. 1C) revealed two major hydrophobic regions (see round-ended boxes in Fig. 1D). One, at its NH₂-terminus, is apparently the signal peptide whose most likely cleavage site lies between the glycine and isoleucine residues designated in Fig. 1D as -1 and +1 respectively. The other major hydrophobic domain, located between residues 191 and 213, is flanked at both ends by several charged amino acids, characteristic of a membrane anchoring
55 domain. As in several other transmembrane proteins, the amino acids confining the hydrophobic domain at its COOH-terminal are basic. The transmembrane domain bisects the predicted protein into almost equally sized extracellular and intracellular domains.

The extracellular domain has 3 putative sites for asparagine-linked glycosylation (overlined in Fig. 1D).

Assuming that the amount of oligosaccharides in the extracellular domain is similar to that reported as present in TBP-I (Seckinger, P., et al., (1989b) Cytokine I, 149 (an abstract)), the molecular size of the mature protein is very similar to that estimated for the type I receptor (about 58kD) (Hohmann, H.P., et al., (1989) J.Biol.Chem. 264, pp. 14927-14934).

5 b) Features of the predicted amino acid sequence in the Type I TNF-R and relationship to the structure of TBP-I and TBP-II

The amino acid sequence of the extracellular domain of the protein encoded by the E13 cDNA fully
 10 matches several sequences of amino acids determined in TBP-I (Table I). It contains the NH₂-terminal amino acid sequence of TBP-I at amino acids 20-32 (compare Fig. 1D and Fig. 1Aa), a sequence corresponding to the COOH terminus of TBP-I at amino acid 178-180; and, also, adjacent to the first methionine located further downstream in the encoded protein, a sequence which is identical to the NH₂-terminal amino acid sequence of a cyanogenbromide cleavage fragment of TBP-I (broken lines in Fig. 1D).
 15 There is also a marked similarity in amino acid composition between the extracellular domain of the receptor and TBP-I (Table II).

The most salient feature of this amino acid composition is a very high content of cystein residues (shown boxed in Fig. 1D). The positioning of the cystein residues as well as of other amino acids within the extracellular domain displays a four-fold repetition pattern (underlined in Fig. 1D). The amino acid sequence
 20 within the extracellular domain of the TNF-R, which corresponds to the COOH terminal sequence of TBP-I (see Table I and Fig. 4), is located at the COOH terminus of the cystein-rich repeat region. The sequence corresponding to the NH₂ terminal sequence of TBP-I corresponds to a sequence located a few amino acids upstream of the NH₂ terminal end of this region (broken lines in Fig. 1D) in the extracellular domain.

In contrast to the identity of amino acid sequences between TBP-I and the extracellular domain of the
 25 type I TNF receptor, sequences examined in the soluble form of the type II TNF-R (TBP-II, Table * I) were not identical to any sequence in the type I TNF-R. This finding is expected, considering the lack of immunological crossreactivity between the two receptors (Engelmann, H., et al., (1990) J.Biol.Chem. 265, pp. 1531-1536).

In contrast to the very high content of cystein residues in the putative extracellular domain of the type I
 30 TNF-R, there are only 5 cystein residues in the intracellular domain. Between the two which are proximal to the transmembrane domain (positions 227 and 283), extends a stretch of 55 amino acids which is rich in proline residues (15% of the residues) and even richer in serine and threonine residues (36%), most located very close to or adjacent to each other.

35 EXAMPLE 3

Expression of the type I TNF-R cDNA

To explore the relation between the protein encoded by the E13 cDNA and TBP-I further, this protein
 40 was expressed in CHO cells. The E13 cDNA was introduced into an expression vector and was cotransfected with a recombinant vector containing the dihydrofolate reductase (DHFR) cDNA into DHFR-deficient cells. After selection by growth in a nucleotide-free medium, individual clones were amplified by growth in the presence of methotrexate. A number of clones which react with several monoclonal antibodies that bind to spatially distinct epitopes in TBP-I were detected (Fig.2). Expression of the protein was correlated with an
 45 increase in specific binding of human TNF to the cells (Table III).

Applying a sensitive immunoassay for TBP-I in which polyclonal antibodies and a monoclonal antibody against this protein were employed, (Procedures, Example 1f) in the medium of CHO cells which express the human TNF-R on their surface, also a soluble form of the protein could be detected (Table III). All of five
 50 different CHO clones which expressed the TNF-R, produced this soluble protein. Several other transfected clones which did not express the cell surface receptor did not produce its soluble form either. When analyzed by reversed phase HPLC, the CHO-produced soluble TNF-R eluted as a single peak, with a retention time identical to that of TBP-I (Fig. 3).

EXAMPLE 4

Cloning of the cDNA encoding TBP-I and expression of TBP-I in Chinese Hamster Ovary (CHO) cells

In order to obtain plasmids suitable for efficient expression of the DNA encoding a soluble domain of

the type I TNF receptor in mammalian cells, the gene from position 256 to position 858 of the DNA sequence shown in Fig. 1D, was cloned in two expression vectors: in one plasmid, gene expression was under the SV40 early gene promoter; in the second plasmid, gene expression was under the regulation of the cytomegalovirus (CMV) promoter. These vectors were introduced into CHO cells by CaPO₄ coprecipitation with a plasmid DHFR selectable genetic marker.

Construction of Expression Vectors

1) SV40 Early Promoter-TBP-I fusion: Plasmid pSV-TBP.

Constitutive expression of TBP-I can be achieved by using an expression vector that contains the DNA sequence coding for TBP-I fused to the strong SV40 early gene promoter (Fig 5).

Step 1: A DNA fragment coding for TBP-I, including its signal peptide and extending to amino acid 180 was prepared by PCR amplification. For amplification two oligonucleotides were used as primers: the 5' end primer contains the sequence coding for the first seven amino acids of the signal peptide, preceded by six nucleotides; the 3' end oligonucleotide contains the sequence coding for amino acid residues 174 through 180 followed by two stop codons (TGA and TAA).

The conditions for PCR amplification are the following:

	Temperature °C	Time min
1 cycle	94	6
	50	2
	72	4
30 cycles	94	1
	50	2
	72	4
1 cycle	94	1
	50	2
	72	12

Step 2: After sequence verification, the amplified DNA fragment was cloned into the HincII restriction site of plasmid pGEM-1 by blunt end ligation. Plasmids pTBP-16 and pTBP-17 are the two plasmids obtained in this way and they differ in the orientation of the TBP-I insert with respect to the cloning vector.

Step 3: The DNA fragment containing TBP-I was excised from plasmid pTBP-17 using the two adjacent restriction sites HindIII (at the 5' end) and BamHI (at the 3' end).

Step 4: Finally, this fragment was cloned between the HindIII and the BclI restriction sites of the expression vector pSVE3.

The resulting plasmid is called pSV-TBP (Fig. 5).

2) CMV promoter-TBP-I fusion: plasmid pCMV-TBP.

Alternatively, constitutive expression of TBP-I can be achieved by using an expression vector that contains the DNA sequence coding for TBP-I fused to the CMV promoter (Fig 6).

The first two steps for the construction of the CMV based vector are identical to the ones described for the construction of the SV40-TBFI fusion plasmid, as described above.

Step 3: The DNA fragment containing TBP-I was excised from plasmid pTBP-17 using the two adjacent restriction sites HindIII (at the 5' end) and XbaI (at the 3' end).

Step 4: Finally, this fragment was cloned between the HindIII and the XbaI restriction sites of the expression vector Rc/CMV.

The resulting plasmid is called pCMV-TBP.

Expression of Human TBP-I in CHO Cells

5 CHO cells CHO-K1 DHFR⁻, lacking DHFR activity, were transformed by CaPO₄ coprecipitation with a 12:1 mixture of uncut pSV-TBP DNA (73 µg) and mpSV2DHFR (6 µg) DNA, the latter being the selectable marker. Alternatively, CHO cells were transformed with a 5:1 mixture of pCMV-TBP (30 µg) and mpSV2DHFR (6 µg).

Cells were grown in nutrient mixture F12 (Gibco) with 10% fetal calf serum (FCS) at 37° C in 5% CO₂.
 10 For DNA transfection, 5x10⁵ cells were cultured for one day in 9 cm plates. A CaPO₄-DNA coprecipitate was prepared by mixing plasmid DNAs, dissolved in 0.45 ml of Tris-HCl pH 7.9, 0.1 mM EDTA with 0.05 ml of 2.5 M CaCl₂; thereafter, 0.5 ml of 280 mM Na₂PO₄, 50 mM Hepes buffer pH 7.1 was added with gentle mixing. The mixture was kept for 30-40 minutes at room temperature in order to form the precipitate. After adding the CaPO₄-DNA to the cells and leaving the cells at room temperature for 30 min, 9 ml of nutrient
 15 mixture F12, 10% FCS were added and the cultures returned to the CO₂ incubator for 4 hours. Medium was removed and the cells were osmotically shocked with 10% glycerol in F12 for 4 min. After 48 hours of growth in non-selective medium, the cells were then trypsinized and subcultured 1:10 into selective medium, composed of Dulbecco's modified Eagle's medium (DMEM) (H21, Gibco), 150 µg/ml proline, and 10% FCS which had been extensively dialyzed against phosphate-buffered saline (PBS). In some cases,
 20 MEM alpha medium without nucleotides (F20, Gibco) was used. The cultures were kept at 37° C and 10% CO₂ and the medium was changed every 3-4 days. Clones were isolated after about 15 days, trypsinized, and grown to mass cultures.

Transformants able to grow in medium lacking thymidine (DMEM with dialyzed serum) were obtained. Culture supernatants of individual transformant clones or culture supernatant of mixed populations were
 25 screened for human TBP-I by measuring the level of secreted protein by the enzyme linked immunoassay described in Example 1f. TBP-I levels of up to 10 ng/ml were detected in culture supernatants of mixed cells populations.

This example shown that TBP-I or a similar soluble protein can be obtained also by transfection of mammalian cells with a DNA encoding the soluble domain of the type I TNF receptor.

30

EXAMPLE 5

Expression of TBP-I in E. coli.

35 For expression of TBP-I in E.coli, the sequence coding for the signal peptide and for the first 19 aminoacids (Arg) must be removed (Figure 1D). Moreover, the Asp residue must be preceded by a Met residue. The desired fragment is then cloned into the expression vector pKK223-3 that contains the hybrid tryp-lac promoter. To achieve this goal plasmid pTBP-16 (Fig 5) is cut with the two unique restriction sites StyI and HindIII. StyI restriction site is C/CAAGG and, therefore, it cuts after Pro24. HindIII restriction site is
 40 located in the polylinker of the plasmid and downstream from the two added stop codons that follow Asn180 (Fig. 5).

The resulting DNA fragment, coding for TBP-I, has an intact 3' end and a truncated 5' end, where the sequence preceding the StyI site and coding for Asp-Ser-Val-Cys-Pro has been removed.

For cloning of the StyI-HindIII fragment into the expression vector pKK223-3, the following couple of
 45 synthetic oligonucleotides are used:

		Met Asp Ser Val Cys Pro	
50	5'	AATTC ATG CAT AGT GTG TGT CCC	3'
	3'	G TAC CTA TCA CAC ACA GGG GTT C	5'
		┌──────────┐ ┌──────────┐	
55		EcoRI StyI	

One end of this double stranded oligonucleotide is an EcoRI restriction site. This end is ligated to the EcoRI site of plasmid pKK223-3, located downstream to the tryp-lac promoter. The second end of the double stranded oligonucleotide is a Styl restriction site to be ligated to the Styl of the TBP-I DNA fragment.

The remainder of the sequence is such that the codons coding for the first five amino acids are restored and that an additional Met codon is added in front of Asp20. The expression vector is obtained by ligation of plasmid pKK223-3, digested with EcoRI and HindIII, to the double-stranded synthetic oligonucleotide and to the Styl-HindIII TBPI fragment.

E.coli cells are transfected with this expression vector in order to produce TBP-I.

10

15

20

25

30

35

40

45

50

55

Table I: Amino acid sequences of TBP I and TBP II

TBP I:

NH₂-1 (-N-terminus)NH₂-2

C-terminus

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
NH ₂	Asp	Ser	Val	Cys	Pro	Gln	Gly	Lys	Tyr	Ile	His	Pro	Gln	---
NH ₂	Gly	Gln	Val	Glu	Leu	Ser	Ser	Cys	Thr	Val	Asp	Arg	Thr	Val
---	Ile	Glu	Asn	COOH										---

TBP II:

N-terminus

TRP 35

TRP 38/1

TRP 39/2

TRP 44/1

TRP 44/2

TRP 46/1

TRP 46/2

TRP 50

TRP 54/1

TRP 54/2

TRP 53/1

TRP 53/2

TRP 60

TRP 62

TRP 65

TRP 67

TRP 84

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
NH ₂	Ala	Gln	Val	Ala	Phe	Thr	Tyr	Ala	Pro	Glu	Pro	Gly	Ser	Thr	Cys	Arg	Leu	Arg	Glu	Tyr	Tyr
NH ₂	Leu	Cys	Ala	Pro	Leu	Arg	Lys														
NH ₂	Cys	Arg	Pro	Gly	Phe	Gly	Val	Ala	Arg												
NH ₂	Glu	Tyr	Tyr	Asp	Gln	Thr	Ala	Gln	Met	Cys	Cys	---									
NH ₂	Glu	Tyr	Tyr	Asp	Gln	Thr	Ala	Gln	Met	Cys	Ser	---									
NH ₂	Ser	Cys	Gly	Pro	Ser	Tyr	Pro	Asp	---												
NH ₂	Phe	Thr	Pro	Tyr	Ala	Pro	Glu	Pro	Gly	Thr	Cys	Arg									
NH ₂	Leu	Arg	Glu	Tyr	Tyr	Asp	Gln	Thr	Ala	Gln	Met	Cys	Cys	---							
NH ₂	Leu	Arg	Glu	Tyr	Tyr	Asp	Gln	Thr	Ala	Gln	Met	Cys	Cys	---							
NH ₂	Pro	Gly	Trp	Tyr	Cys	Ala	Leu	Ser	Lys												
NH ₂	Ala	Gln	Val	Ala	Phe	Thr	Pro	Tyr	Ala	Pro	Glu	Pro	Gly	Ser	Thr	Cys	Arg				
NH ₂	Val	Ala	Phe	Thr	Pro	Tyr	Ala	Pro	Glu	Pro	Gly	Ser	Thr	Cys	Arg						
NH ₂	Cys	Arg	Pro	Gly	Phe	Gly	Val	Ala	Arg												
NH ₂	Ile	Cys	Thr	Cys	Arg	Pro	Gly	Trp	Tyr	Cys	Ala	Leu	Ser	---							
NH ₂	Pro	Gly	Thr	Glu	Thr	Ser	Asp	Val	Val	Cys	Lys	Pro	Cys	Ala	Pro	Gly	Thr	Phe	Ser	Lys	
NH ₂	Pro	Gly	Thr	Glu	Thr	Ser	Asp	Val	Val	Cys	Lys	Pro	Cys	Ala	Pro	Gly	Thr	Phe	Ser	Lys	
NH ₂	Pro	Gly	Thr	Glu	Thr	Ser	Asp	Val	Val	Cys	Lys	Pro	Cys	Ala	Pro	Gly	Thr	Phe	Ser	Lys	
NH ₂	Cys	Arg	Pro	Gly	Phe	Gly	Val	Ala	Arg	Pro	Glu	Thr	Glu	Thr	Ser	Asp	Val	Val	Cys	Lys	
NH ₂	Thr	Ser	Asp	Thr	Val	Cys	Asp	Ser	Cys	Glu	Asp	Ser	Thr	Tyr	Thr	Gln	Leu	Trp	---		

Table II. Similarity of the amino acid compositions of the TNF binding protein TBPI and a corresponding region in the extracellular domain of the TNF-R (type I)

5	Amino acid	mol/100 mol of amino acids	
		TBPI*	Residues 20-180 in the extracellular domain**
	Ala 1.7	1.2	
10	Cys	12.8	14.9
	Asp + Asn	10.9	11.1
	Glu + Gln	13.9	12.4
15	Phe	3.2	3.1
	Gly	8.3	5.6
	Ile	4.4	4.3
	Ile	2.8	2.5
20	Lys	6.2	6.2
	Leu	8.0	6.8
	Met	0.4	0.0
25	Pro	3.8	3.1
	Arg	4.7	4.3
	Ser	8.1	9.3
	Thr	6.1	6.2
30	Val	4.2	4.3
	Trp	-	0.6
	Tyr	2.4	3.1

35 * According to Olsson et al., 1989

**Residue 20 corresponds to the NH₂-terminal amino acid of TBPI. Residue 180 is the COOH-terminal residue of TBPI.

40

45

50

55

Table III. Expression of the cell surface and soluble forms of human type I TNF-R in CHO cells

CHO cell clone	Specific binding of TNF (CPM/10 ⁶ cells)	cells expressing human cell surface TNF-R (% fluorescent cells)	human soluble type I TNF receptors (pg/ml)
nontransfected	180±46	<1%	<0.03
C8	175±60	<1%	<0.03
IL-16	650±60	73%	30
IL-18	610±40	80%	40

The IL-16 and IL-18 clones consist of cells transfected with a recombinant expression vector containing B13 cDNA. C-8 cells were transfected with a control vector (see Fig. 3). Binding of radiolabelled TNF to the cells was determined in pentuplicate samples. Detection of immunoreactive receptors on the surface of the cells was carried out using combined 17, 18, 20 and 30 anti TBP-I monoclonal antibodies. Results are expressed as percentage of fluorescent cells (background values, obtained by staining the cells with an anti-TNF monoclonal antibody, are subtracted). For other details, see Materials and Methods.

Claims

1. A method for the production of a soluble recombinant protein selected from human Tumor Necrosis Factor Binding Protein I (TBP-I), biologically active precursors and analogs thereof, which comprises:
 - i) transfecting eukaryotic cells with an expression vector comprising a DNA molecule encoding the whole human type I TNF receptor or a soluble domain thereof, and
 - ii) culturing the transfected cells, whereby the desired protein is produced and secreted into the medium.
2. A method according to claim 1 wherein the DNA molecule encoding the whole type I TNF receptor is the cDNA having the sequence depicted in Figure 1D.
3. A method according to claim 2 wherein the cDNA is introduced into an expression vector and is cotransfected with a recombinant vector containing the dihydrofolate reductase (DHFR) cDNA into DHFR-deficient chinese hamster ovary (CHO) cells.
4. A method according to claim 3 wherein the cells are selected by growth in a nucleotide-free medium, individual clones are amplified by growth in the presence of methotrexate and the soluble protein secreted into the medium is detected by reaction with monoclonal and polyclonal antibodies raised against TBP-I.
5. A method according to any of claims 1 to 4 wherein the soluble protein secreted into the medium shows a retention time identical to that of TBP-I when analyzed by reversed phase HPLC.
6. A method according to any of claims 1 to 5 for the production of human TBP-I.
7. A method according to any of claims 1 to 5 for the production of a human TBP-I precursor or analog.
8. A soluble protein selected from precursors and analogs of TBP-I which are secreted into the medium of eukaryotic cells transfected with a cDNA encoding the whole type I human TNF receptor or a soluble domain thereof.
9. A soluble protein as claimed in claim 8 secreted into the medium of CHO cells transfected with the cDNA molecule depicted in Figure 1D.


A a Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln

b GGI GTC CCI TTC ATA TAA GTA GGI GT
T T G G G T

c GGA GTC CCA TTC ATA TA
C T C C T
G T

d TTC ATA TAA GTA GGA GT
T G G G C
T C T

B



C




Figure 1 A-C

D	18	ACCCCAAGGCCAGACTGGAGCCCTCAGTCCAGAGATTCTGAGAAATTAAGCAGAGAGGAGGAGATCAGTGGACCGCCGTGATCTCTATGCGCCGAGTCTCAACCCCTCAA	17
	137	CTGTCAACCCCAAGGCCAGACTTGGAGAGTCTGGAGACAGCCAGTCCGGGMAAGCCAGCACTGGCCCTGGCCACACTGGCTGAGCCCAATGGGGAGTGAGAGCCCATAGCTGTCTGGC	136
		-21 -10 -1 +1	255
	256	ATG GGC CTC TCC ACC GTG CCT GAC CTG CTG CCG CTG GTG CTG CTG GAG CTG TTG GTG GGA ATA TAC CCC TCA GGC GTT ATT GGA CTG	345
		Met Gly Leu Ser Thr Val Pro Asp Leu Leu Leu Val Leu Leu Val Gly Ile Tyr Pro Ser Gly Val Ile Gly Leu	
	346	Val Pro His Leu Gly Asp Arg Gly Lys Arg Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Ser Ile Cys Cys Thr	435
		GTC CCT CAC CTA GGG GAC AGG GAG AAG AGA GAT AGT GTG TGT CCC CAA GGA AAA TAT ATC CAC CCT CAA AAT TCG ATT TGC TGT ACC	
	436	Lys Cys His Lys Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr	525
		AAG TGC CAC AAA GGA ACC TAC TTG TAC AAT GAC TGT CCA GGC CCG GGC CAG GAT ACG GAC TGC AGG GAG TGT GAG AGC GGC TCC TTC ACC	
	526	Ala Ser Glu Asn His Leu Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val Ile Ser Ser Cys Thr Val Asp	615
		GCT TCA GAA AAC CAC CTC AGA CAC TGC CTC AGC TGC TCC AAA TGC CGA AAG GAA ATG GGT CAG GTG GAG ATC TCT TGC ACA GTG GAC	
	616	Arg Asp Thr Val Cys Gly Arg Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Cys Phe Asn Cys Ser Leu Cys Leu	705
		CGG GAC ACC GTG TGT GGC TGC AGG AAG AAC CAG TAC CCG CAT TAT TGG AGT GAA AAC CTT TTC CAG TGC TTC AAT TGC AGC CTC TGC CTC	
	706	Asn Gly Thr Val His Leu Ser Cys Gln Glu Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu Asn Glu Cys Val	795
		AAT GGG ACC CTG CAC CTC TCC TGC CAG GAG AAA CAG AAC ACC GTG TGC ACC TGC CAT GCA GGT TTC TTT CTA AGA GAA AAC GAG TGT GTC	
	796	Ser Cys Ser Asn Cys Lys Lys Ser Leu Glu Cys Thr Lys Leu Cys Leu Pro Gln Ile Glu Asn Val Lys Gly Thr Glu Asp Ser Gly Thr	885
		TCC TGT AGT AAC TGT AAG AAA AGC CTG GAG TGC ACG AAG TTG TGC CTA CCC CAG ATT GAG AAT GTT AAG GGC ACT GAG GAC TCA GGC ACC	
	886	Thr Val Leu Leu pro Leu Val Ile Phe Phe Gly Leu Cys Leu Leu Ser Leu Leu Phe Ile Gly Leu Met Tyr Arg Tyr Gln Arg Tyr Lys	975
		ACA GTG CTG TTG CCC CTG GTC ATT TTC TTT GGT CTT TGC CTT TTA TCC CTC CTC TTC ATT GGT TTA ATG TAT CCG TAC CAA CGG TGG AAG	

Figure 1 D (part 1)

220 Ser Lys Leu Tyr Ser Ile Val [Cys] Gly Lys Ser Thr Pro Glu Lys Glu Gly Glu Leu Glu Gly Thr Thr Lys Pro Leu Ala Pro Asn
 230 TCC AAG CTC TAC TCC ATT GTT TGT GCG AAA TCG ACA CCT GAA AAA GAG GCG GAG CTT GAA GCA ACT ACT ACT AAG CCC CTG GCC CCA AAC
 240
 250
 260
 270
 280
 290
 300
 310
 320
 330
 340
 350
 360
 370
 380
 390
 400
 410
 420
 430
 440
 450
 460
 470
 480
 490
 500
 510
 520
 530
 540
 550
 560
 570
 580
 590
 600
 610
 620
 630
 640
 650
 660
 670
 680
 690
 700
 710
 720
 730
 740
 750
 760
 770
 780
 790
 800
 810
 820
 830
 840
 850
 860
 870
 880
 890
 900
 910
 920
 930
 940
 950
 960
 970
 980
 990
 1000
 1010
 1020
 1030
 1040
 1050
 1060
 1070
 1080
 1090
 1100
 1110
 1120
 1130
 1140
 1150
 1160
 1170
 1180
 1190
 1200
 1210
 1220
 1230
 1240
 1250
 1260
 1270
 1280
 1290
 1300
 1310
 1320
 1330
 1340
 1350
 1360
 1370
 1380
 1390
 1400
 1410
 1420
 1430
 1440
 1450
 1460
 1470
 1480
 1490
 1500
 1510
 1520
 1530
 1540
 1550
 1560
 1570
 1580
 1590
 1600
 1610
 1620
 1630
 1640
 1650
 1660
 1670
 1680
 1690
 1700
 1710
 1720
 1730
 1740
 1750
 1760
 1770
 1780
 1790
 1800
 1810
 1820
 1830
 1840
 1850
 1860
 1870
 1880
 1890
 1900
 1910
 1920
 1930
 1940
 1950
 1960
 1970
 1980
 1990
 2000
 2010
 2020
 2030
 2040
 2050
 2060
 2070
 2080
 2090
 2100
 2110
 2120
 2130
 2140
 2150
 2160
 2170
 2180
 2190
 2200
 2210
 2220
 2230
 2240
 2250
 2260
 2270
 2280
 2290
 2300
 2310
 2320
 2330
 2340
 2350
 2360
 2370
 2380
 2390
 2400
 2410
 2420
 2430
 2440
 2450
 2460
 2470
 2480
 2490
 2500
 2510
 2520
 2530
 2540
 2550
 2560
 2570
 2580
 2590
 2600
 2610
 2620
 2630
 2640
 2650
 2660
 2670
 2680
 2690
 2700
 2710
 2720
 2730
 2740
 2750
 2760
 2770
 2780
 2790
 2800
 2810
 2820
 2830
 2840
 2850
 2860
 2870
 2880
 2890
 2900
 2910
 2920
 2930
 2940
 2950
 2960
 2970
 2980
 2990
 3000
 3010
 3020
 3030
 3040
 3050
 3060
 3070
 3080
 3090
 3100
 3110
 3120
 3130
 3140
 3150
 3160
 3170
 3180
 3190
 3200
 3210
 3220
 3230
 3240
 3250
 3260
 3270
 3280
 3290
 3300
 3310
 3320
 3330
 3340
 3350
 3360
 3370
 3380
 3390
 3400
 3410
 3420
 3430
 3440
 3450
 3460
 3470
 3480
 3490
 3500
 3510
 3520
 3530
 3540
 3550
 3560
 3570
 3580
 3590
 3600
 3610
 3620
 3630
 3640
 3650
 3660
 3670
 3680
 3690
 3700
 3710
 3720
 3730
 3740
 3750
 3760
 3770
 3780
 3790
 3800
 3810
 3820
 3830
 3840
 3850
 3860
 3870
 3880
 3890
 3900
 3910
 3920
 3930
 3940
 3950
 3960
 3970
 3980
 3990
 4000
 4010
 4020
 4030
 4040
 4050
 4060
 4070
 4080
 4090
 4100
 4110
 4120
 4130
 4140
 4150
 4160
 4170
 4180
 4190
 4200
 4210
 4220
 4230
 4240
 4250
 4260
 4270
 4280
 4290
 4300
 4310
 4320
 4330
 4340
 4350
 4360
 4370
 4380
 4390
 4400
 4410
 4420
 4430
 4440
 4450
 4460
 4470
 4480
 4490
 4500
 4510
 4520
 4530
 4540
 4550
 4560
 4570
 4580
 4590
 4600
 4610
 4620
 4630
 4640
 4650
 4660
 4670
 4680
 4690
 4700
 4710
 4720
 4730
 4740
 4750
 4760
 4770
 4780
 4790
 4800
 4810
 4820
 4830
 4840
 4850
 4860
 4870
 4880
 4890
 4900
 4910
 4920
 4930
 4940
 4950
 4960
 4970
 4980
 4990
 5000
 5010
 5020
 5030
 5040
 5050
 5060
 5070
 5080
 5090
 5100
 5110
 5120
 5130
 5140
 5150
 5160
 5170
 5180
 5190
 5200
 5210
 5220
 5230
 5240
 5250
 5260
 5270
 5280
 5290
 5300
 5310
 5320
 5330
 5340
 5350
 5360
 5370
 5380
 5390
 5400
 5410
 5420
 5430
 5440
 5450
 5460
 5470
 5480
 5490
 5500
 5510
 5520
 5530
 5540
 5550
 5560
 5570
 5580
 5590
 5600
 5610
 5620
 5630
 5640
 5650
 5660
 5670
 5680
 5690
 5700
 5710
 5720
 5730
 5740
 5750
 5760
 5770
 5780
 5790
 5800
 5810
 5820
 5830
 5840
 5850
 5860
 5870
 5880
 5890
 5900
 5910
 5920
 5930
 5940
 5950
 5960
 5970
 5980
 5990
 6000
 6010
 6020
 6030
 6040
 6050
 6060
 6070
 6080
 6090
 6100
 6110
 6120
 6130
 6140
 6150
 6160
 6170
 6180
 6190
 6200
 6210
 6220
 6230
 6240
 6250
 6260
 6270
 6280
 6290
 6300
 6310
 6320
 6330
 6340
 6350
 6360
 6370
 6380
 6390
 6400
 6410
 6420
 6430
 6440
 6450
 6460
 6470
 6480
 6490
 6500
 6510
 6520
 6530
 6540
 6550
 6560
 6570
 6580
 6590
 6600
 6610
 6620
 6630
 6640
 6650
 6660
 6670
 6680
 6690
 6700
 6710
 6720
 6730
 6740
 6750
 6760
 6770
 6780
 6790
 6800
 6810
 6820
 6830
 6840
 6850
 6860
 6870
 6880
 6890
 6900
 6910
 6920
 6930
 6940
 6950
 6960
 6970
 6980
 6990
 7000
 7010
 7020
 7030
 7040
 7050
 7060
 7070
 7080
 7090
 7100
 7110
 7120
 7130
 7140
 7150
 7160
 7170
 7180
 7190
 7200
 7210
 7220
 7230
 7240
 7250
 7260
 7270
 7280
 7290
 7300
 7310
 7320
 7330
 7340
 7350
 7360
 7370
 7380
 7390
 7400
 7410
 7420
 7430
 7440
 7450
 7460
 7470
 7480
 7490
 7500
 7510
 7520
 7530
 7540
 7550
 7560
 7570
 7580
 7590
 7600
 7610
 7620
 7630
 7640
 7650
 7660
 7670
 7680
 7690
 7700
 7710
 7720
 7730
 7740
 7750
 7760
 7770
 7780
 7790
 7800
 7810
 7820
 7830
 7840
 7850
 7860
 7870
 7880
 7890
 7900
 7910
 7920
 7930
 7940
 7950
 7960
 7970
 7980
 7990
 8000
 8010
 8020
 8030
 8040
 8050
 8060
 8070
 8080
 8090
 8100
 8110
 8120
 8130
 8140
 8150
 8160
 8170
 8180
 8190
 8200
 8210
 8220
 8230
 8240
 8250
 8260
 8270
 8280
 8290
 8300
 8310
 8320
 8330
 8340
 8350
 8360
 8370
 8380
 8390
 8400
 8410
 8420
 8430
 8440
 8450
 8460
 8470
 8480
 8490
 8500
 8510
 8520
 8530
 8540
 8550
 8560
 8570
 8580
 8590
 8600
 8610
 8620
 8630
 8640
 8650
 8660
 8670
 8680
 8690
 8700
 8710
 8720
 8730
 8740
 8750
 8760
 8770
 8780
 8790
 8800
 8810
 8820
 8830
 8840
 8850
 8860
 8870
 8880
 8890
 8900
 8910
 8920
 8930
 8940
 8950
 8960
 8970
 8980
 8990
 9000
 9010
 9020
 9030
 9040
 9050
 9060
 9070
 9080
 9090
 9100
 9110
 9120
 9130
 9140
 9150
 9160
 9170
 9180
 9190
 9200
 9210
 9220
 9230
 9240
 9250
 9260
 9270
 9280
 9290
 9300
 9310
 9320
 9330
 9340
 9350
 9360
 9370
 9380
 9390
 9400
 9410
 9420
 9430
 9440
 9450
 9460
 9470
 9480
 9490
 9500
 9510
 9520
 9530
 9540
 9550
 9560
 9570
 9580
 9590
 9600
 9610
 9620
 9630
 9640
 9650
 9660
 9670
 9680
 9690
 9700
 9710
 9720
 9730
 9740
 9750
 9760
 9770
 9780
 9790
 9800
 9810
 9820
 9830
 9840
 9850
 9860
 9870
 9880
 9890
 9900
 9910
 9920
 9930
 9940
 9950
 9960
 9970
 9980
 9990
 10000
 10010
 10020
 10030
 10040
 10050
 10060
 10070
 10080
 10090
 10100
 10110
 10120
 10130
 10140
 10150
 10160
 10170
 10180
 10190
 10200
 10210
 10220
 10230
 10240
 10250
 10260
 10270
 10280
 10290
 10300
 10310
 10320
 10330
 10340
 10350
 10360
 10370
 10380
 10390
 10400
 10410
 10420
 10430
 10440
 10450
 10460
 10470
 10480
 10490
 10500
 10510
 10520
 10530
 10540
 10550
 10560
 10570
 10580
 10590
 10600
 10610
 10620
 10630
 10640
 10650
 10660
 10670
 10680
 10690
 10700
 10710
 10720
 10730
 10740
 10750
 10760
 10770
 10780
 10790
 10800
 10810
 10820
 10830
 10840
 10850
 10860
 10870
 10880
 10890
 10900
 10910
 10920
 10930
 10940
 10950
 10960
 10970
 10980
 10990
 11000
 11010
 11020
 11030
 11040
 11050
 11060
 11070
 11080
 11090
 11100
 11110
 11120
 11130
 11140
 11150
 11160
 11170
 11180
 11190
 11200
 11210
 11220
 11230
 11240
 11250
 11260
 11270
 11280
 11290
 11300
 11310
 11320
 11330
 11340
 11350
 11360
 11370
 11380
 11390
 11400
 11410
 11420
 11430
 11440
 11450
 11460
 11470
 11480
 11490
 11500
 11510
 11520
 11530
 11540
 11550
 11560
 11570
 11580
 11590
 11600
 11610
 11620
 11630
 11640
 11650
 11660
 11670
 11680
 11690
 11700
 11710
 11720
 11730
 11740
 11750
 11760
 11770
 11780
 11790
 11800
 11810
 11820
 11830
 11840
 11850
 11860
 11870
 11880
 11890
 11900
 11910
 11920
 11930
 11940
 11950
 11960
 11970
 11980
 11990
 12000
 12010
 12020
 12030
 12040
 12050
 12060
 12070
 12080
 12090
 12100
 12110
 12120
 12130
 12140
 12150
 12160
 12170
 12180
 12190
 12200
 12210
 12220
 12230
 12240
 12250
 12260
 12270
 12280
 12290
 12300
 12310
 12320
 12330
 12340
 12350
 12360
 12370
 12380
 12390
 12400
 12410
 12420
 12430
 12440
 12450
 12460
 12470
 12480
 12490
 12500
 12510
 12520
 12530
 12540
 12550
 12560
 12570
 12580
 12590
 12600
 12610
 12620
 12630
 12640
 12650
 12660
 12670
 12680
 12690
 12700
 12710
 12720
 12730
 12740
 12750
 12760
 12770
 12780
 12790
 12800
 12810
 12820
 12830
 12840
 12850
 12860
 12870
 12880
 12890
 12900
 12910
 12920
 12930
 12940
 12950
 12960
 12970
 12980
 12990
 13000
 13010
 13020
 13030
 13040
 13050
 13060
 13070
 13080
 13090
 13100
 13110
 13120
 13130
 13140
 13150
 13160
 13170
 13180
 13190
 13200
 13210
 13220
 13230
 13240
 13250
 13260
 13270
 13280
 13290
 13300
 13310
 13320
 13330
 13340
 13350
 13360
 13370
 13380
 13390
 13400
 13410
 13420
 13430
 13440
 13450
 13460
 13470
 13480
 134

430
Pro Ser Leu Leu Arg End
1606 CCC AGT CTT CTC AGA TGA GGCTGGCGCCCTGCGGCAGCTCTAAGACCGTCTCGAGATCGCCTTCCACCCACCTTTTCTGAAAGGAGGGTCTCTGCAGGGCAAGC 1718
1719 AGGAGCTAGCAGCGGCTACTTTGGTCTAACCCTCGATGTATAGCTTTTCTCAGCTGCTGCGCGCGCCGACAGTCAGCGCTGTGCGCGGAGAGAGGTGCGCGTGGCTCAA 1837
1838 GAGCTGAGTGGTGGTTTGGGAGGATGAGGACGCTATGCCCTCATGCCGTTTTTGGGTGCTCTCACAGCAAGGCTGCTCGGGGGCCCTGTTTCGTCCCTGAGCCCTTTTTCACAGTG 1956
1957 CATAGCAGTTTTTTTTTTTTTTTTTTTTTTTTTTTAAATCAATCAATGTTACACTAATAGAACTTGGCACTCCTGTGCCCCTCTGCTGGACACATAGCAAGCTGAA 2075
2076 CTGTCTAAGGCAGGCGGAGCAGGACATGGGGCTTCAGCTGGAGCTGTGGACTTTTGTATACACTAAATTTCTGAGTTTAAAAA 2176

Figure 1 D (part 3)




Figure 2




Figure 3




Figure 4

CONSTRUCTION OF PLASMID pSV-TBP






Figure 5 part 1)



. Figure 5 (part 2)




Figure 5 (part 3)

CONSTRUCTION OF PLASMID pCMV-TBP





Figure 6 (part 1)




Figure 6 (part 2)




Figure 6 (part 3)




Figure 6 (part 4)



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 12 4133

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
D,Y	EUROPEAN JOURNAL OF HAEMATOLOGY, vol. 42, no. 3, March 1989, pages 270-275; I. OLSSON et al.: "Isolation and characterization of a tumor necrosis factor binding protein from urine" * Whole document *	1-3,5-9	C 12 N 15/12 C 07 K 13/00 C 12 N 15/79
Y	THE EMBO JOURNAL, vol. 8, no. 9, September 1989, pages 2497-2502; C. BINKERT et al.: "Cloning, sequence analysis and expression of a cDNA encoding a novel insulin-like growth factor binding protein (IGFBP-2)" * Summary *	1-3,5-9	
P,X	CELL, vol. 61, 20th April 1990, pages 351-359, Cambridge, MA, US; H.R. LOETSCHER et al.: "Molecular cloning and expression of the human 55kd tumor necrosis factor receptor" * Summary; figure 2; Experimental procedures *	1,2,5-9	
P,X	PROC. NATL. ACAD. SCI. USA, vol. 87, 14th November 1990, pages 8781-8784, US; T. GATANAGA et al.: "Purification and characterization of an inhibitor (soluble tumor necrosis factor receptor) for tumor necrosis factor and lymphotoxin obtained from the serum ultrafiltrates of human cancer patients" * Whole document *	1,2,5	
P,X	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 265, no. 10, 5th April 1990, pages 5708-5717, US; R.A. HELLER et al.: "Amplified expression of tumor necrosis factor receptor in cells transfected with Epstein-Barr virus shuttle vector cDNA libraries" * Abstract; page 5173, column 2, paragraph 2; page 5174, column 1, paragraph 1: "Discussion" *	1-3,6-9	
The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
The Hague		27 February 91	NAUCHE S.A.
CATEGORY OF CITED DOCUMENTS			
X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention		E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document	



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 12 4133

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	<p>EUROPEAN JOURNAL OF HAEMATOLOGY, vol. 41, no. 5, November 1988, pages 414-419; C. PEETRE et al.: "A tumor necrosis factor binding protein is present in biological fluids"</p> <p>* Whole document *</p> <p>-----</p>	1-9	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
The Hague		27 February 91	NAUCHE S.A.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone</p> <p>Y: particularly relevant if combined with another document of the same category</p> <p>A: technological background</p> <p>O: non-written disclosure</p> <p>P: intermediate document</p> <p>T: theory or principle underlying the invention</p> <p>E: earlier patent document, but published on, or after the filing date</p> <p>D: document cited in the application</p> <p>L: document cited for other reasons</p> <p>S: member of the same patent family, corresponding document</p>			